

## FORMULA BANK THERMODYNAMICS

- Relation between different scales of temperature
 
$$\frac{T_C - 0}{100} = \frac{T_F - 32}{180} = \frac{T_R - 0}{80} = \frac{T_K - 273.15}{100}$$
- $T_C = \frac{5}{9}(T_F - 32)$
- $T_F = \frac{9}{5}T_C + 32$
- $-40^\circ\text{C}$  has same value on Celcius and Fahrenheit scales.
- Triple point of water on absolute scale of temperature is  $273.16\text{ K}$ .
- Faulty Thermometer.
 
$$\frac{\text{False reading} - \text{lower point}}{\text{range}} = \frac{\text{True reading} - \text{lower point}}{\text{range}}$$
- Co-efficient of linear expansion
 
$$\alpha = \frac{\Delta l}{l\Delta T}$$

$$l' = l(1 + \alpha\Delta T)$$
- Coefficient of superficial expansion
 
$$\beta = \frac{\Delta S}{S\Delta T}$$

$$S' = S(1 + \beta\Delta T).$$
- Coefficient of cubic expansion
 
$$\gamma = \frac{\Delta V}{V\Delta T}$$

$$V' = V(1 + \gamma\Delta T).$$
- Relation between  $\alpha$ ,  $\beta$  and  $\gamma$ 

$$6\alpha = 3\beta = 2\gamma$$
 or  $\alpha = \frac{\beta}{2} = \frac{\gamma}{3}$
- Heat supplied to a solid of mass  $m$  for increasing temperature  $\Delta T$  is  $Q = mC\Delta T$ .
- Heat supplied to change its state at constant temperature  $Q = mL$ .
- Gases possess infinite values of specific heat but we consider only two specific heats  $C_p$  and  $C_v$ .
- Mayer's formula  $C_p - C_v = R$ .
- For monoatomic gas,  $f = 3$ 

$$C_v = \frac{3}{2}R \text{ and } C_p = \frac{5}{2}R \text{ and } \gamma = \frac{5}{3} = 1.67$$
- For diatomic gas  $f = 5$  at room temperature
 
$$C_v = \frac{5}{2}R \text{ and } C_p = \frac{7}{2}R \text{ and } \gamma = \frac{7}{5} = 1.4$$
- For triatomic gas  $f = 6$ 

$$C_v = 3R, C_p = 4R \text{ and } \gamma = \frac{4}{3} = 1.33$$
- Joules mechanical equivalent of heat
 
$$J = \frac{W}{Q} = 4.186 \text{ J cal}^{-1}.$$
- Rise in temperature of body when it falls through height  $h$ 

$$\Delta T = \frac{gh}{CJ}$$
- The height from which a block of ice be dropped that it melts completely on reaching ground.
 
$$h = \frac{JL}{g}$$
- The velocity with which a ball of ice be thrown against a wall so that it melts completely,
 
$$v = \sqrt{2JL}$$
- Equation of isothermal process
 
$$PV = \text{Const.}$$
- Equation of adiabatic process
  - $PV^\gamma = \text{Const.}$
  - $TP^{\gamma-1} = \text{Const.}$
  - $\frac{T^\gamma}{P^{\gamma-1}} = \text{Const.}$
- Work done during isothermal process
 
$$W = 2.303 RT \log_{10} \frac{V_2}{V_1}$$

$$W = 2.303 RT \log_{10} \frac{P_2}{P_1}$$
- Work done during adiabatic process
 
$$W = \frac{R}{\gamma-1}(T_1 - T_2)$$

$$W = \frac{R}{\gamma-1}(P_1V_1 - P_2V_2)$$

$$W = C_v(T_1 - T_2)$$
- Slope of adiabatic graph is  $\gamma$ -times more than slope of isothermal process.
- First law of thermodynamics
 
$$dQ = dU + dW$$
- Efficiency of heat engine
 
$$\eta = 1 - \frac{Q_2}{Q_1}$$

$$\eta = 1 - \frac{T_2}{T_1}$$
- Efficiency of heat engine can never be 100%.
- Coefficient of performance of refrigerator.
 
$$\beta = \frac{T_2}{T_1 - T_2} = \frac{Q_2}{Q_1 - Q_2}$$
- There are two dead centres per cycle for a steam engine.